

2021

PHYSICS — HONOURS

Fifth Paper

Full Marks : 100

The figures in the margin indicate full marks.

Candidates are required to give their answers in their own words as far as practicable.

1. Answer **any ten** questions : 2×10
- (a) Is the constraint given by $x\dot{x} + y\dot{y} + x\dot{y} + \dot{x}y = k$ (a constant), a holonomic constraint?
- (b) Show that the two Lagrangians $L_1 = (q - \dot{q})^2$ and $L_2 = (q^2 + \dot{q}^2)$ are equivalent.
- (c) Prove that for motion of a particle under central force, the areal velocity with respect to the centre of force remains constant.
- (d) If the kinetic energy $T = \frac{1}{2} m \dot{r}^2$ and the potential energy $V = \frac{1}{r} \left(1 + \frac{r^2}{c^2} \right)$, find the Hamiltonian 'H' and determine whether $H = T + V$.
- (e) Explain what is meant by streamlines.
- (f) Derive the equation of continuity for a compressible fluid.
- (g) For a four vector A^μ show that $A_\mu A^\mu$ is a scalar.
- (h) Find the constant C which makes $e^{-\alpha x^2}$ an eigenstate of the operator. $\frac{d^2}{dx^2} - Ex^2$ (α is a constant).
- (i) Can we measure the kinetic and potential energies of a particle simultaneously with arbitrary precision?
- (j) Why are the Stokes lines brighter than anti-Stokes lines in Raman Spectra?
- (k) The electronic configuration of Mg is $1s^2 2s^2 2p^6 3s^2$. Obtain its spectral term.
- (l) Why is pure vibrational spectra observed in liquid?

Please Turn Over

Group - A**Section - I****(Classical Mechanics II)**Answer *any two* questions.

2. (a) Starting from Lagrange's equation of motion, obtain Hamilton's equation of motion using Legendre transformation.
- (b) For the Hamiltonian $H = q_1 p_1 - q_2 p_2 - a q_1^2 + b q_2^2$, solve the Hamilton's equation of motion and prove that $q_1 q_2 = \text{constant}$ and $\frac{(p_2 - b q_2)}{q_1} = \text{constant}$.

- (c) Show that the effective potential of a particle of mass 'm' in a central force field is given by

$$U_{\text{eff}}(r) = U(r) + \frac{L^2}{2mr^2}, \text{ where } L \text{ is the angular momentum.} \quad 4+3+3$$

3. (a) Consider a simple harmonic oscillator with angular frequency ω_0 . What will be its angular frequency when a constant force K is applied on it?
- (b) The point of suspension of a simple pendulum moves simple harmonically along the vertical line. Obtain the Lagrangian of the system.
- (c) Prove that, if the Lagrangian of an unconstrained system is invariant under continuous translation, then the total linear momentum is conserved. 3+4+3
4. (a) State Bernoulli's equation of fluid motion and mention the conditions of its validity.
- (b) The Lagrangian of a particle of mass m is $L = \frac{1}{2}(m\dot{x}^2 - bx^2) e^{at}$ where a and b are positive constants. Determine the Hamiltonian. Is it a constant of motion?
- (c) A flat vertical plate is struck normally by a horizontal jet of water 50 mm in diameter with a velocity of 18 m/s. Calculate the force on the plate assuming it to be stationary. 3+4+3

Section - II**(Special Theory of Relativity)**Answer *any two* questions.

5. (a) Define the interval between two events in space time. Show that it is invariant under a Lorentz transformation. Hence explain the conditions for which the interval is time-like, space-like or light-like.
- (b) A muon at rest has life time 2×10^{-6} sec. What is its life time when it travels with a velocity $\frac{3}{5}c$?
- (c) Define covariant and contravariant vector. (1+2+3)+2+2

6. (a) Discuss about inconsistency, if any, in Newton's law of gravitation in the light of postulates of special theory of relativity.
- (b) Define Minkowski space. Show that Lorentz transformation can be regarded as transformation due to a rotation of axes through an imaginary angle given by $\theta = \tan^{-1}(i\beta)$ where $\beta = \frac{v}{c}$ in the 4-dimensional Minkowski space.
- (c) Two rods of proper length l_0 move lengthwise towards each other parallel to the common axis with the same velocity v relative to the laboratory frame. Show that the length of each rod in the reference frame fixed to the other rod is $l = l_0 \frac{(1-\beta^2)}{(1+\beta^2)}$, $\beta = \frac{v}{c}$. 2+(1+3)+4
7. (a) Define proper time interval $d\tau$. Hence construct velocity four vector. Show that it is a time-like vector.
- (b) If $A^{\mu\nu}$ and $B^{\mu\nu}$ are two tensors, Show that $A^{\mu\nu} B_{\mu\nu} = A_{\mu\nu} B^{\mu\nu}$.
- (c) For two four vectors A and B , prove that $A_{\mu} B^{\mu} = A^{\mu} B_{\mu}$. 4+4+2

Group - B

Section - I

(Quantum Mechanics II)

Answer *any two* questions.

8. (a) Consider a one-dimensional simple harmonic oscillator moving in a potential $V(x) = \frac{1}{2}m\omega^2 x^2$.
Given that the ground state wave function is $\psi(x) = \left(\frac{\alpha}{\pi}\right)^{1/4} \exp\left(-\frac{1}{2}\alpha x^2\right)$ (where $\alpha = m\omega/\hbar$).
Find the expectation value of (x^2) .
- (b) For a Hamiltonian $\hat{H} = (\hat{p}^2/2m) + V(\hat{x})$, prove that $[\hat{x}, [\hat{x}, \hat{H}]] = -\frac{\hbar^2}{m}$.
- (c) Prove that $\exp[i(\hat{A}\hat{B} - \hat{B}\hat{A})]$ is a Hermitian operator, if \hat{A}, \hat{B} are Hermitian operators. 4+3+3
9. (a) A stream of particles of mass m and energy E move towards the potential step $V(x) = 0$ for $x < 0$ and $V(x) = V_0$ for $x \geq 0$. If the energy of the particles $E < V_0$,
- (i) show that there is a finite probability of finding the particles in the region $x > 0$.
 - (ii) sketch the solutions in the two regions.
 - (iii) determine the reflection coefficient and comment on the result.
- (b) Write down Pauli's spin matrices σ_x, σ_y and σ_z . The eigenfunctions of the Pauli spin operator σ_z are α and β . Show that $\frac{\alpha + \beta}{\sqrt{2}}$ and $\frac{\alpha - \beta}{\sqrt{2}}$ are the eigenfunctions of σ_x . (3+1+2)+(2+2)

Please Turn Over

10. (a) Write down the Schrödinger equation for the hydrogen atom assuming the nucleus heavy. Obtain the radial part of the equation.
- (b) In the ground state of hydrogen atom show that the probability P for the electron to lie within a sphere of radius R is

$$P = 1 - \exp\left(-\frac{2R}{a_0}\right) \left(1 + \frac{2R}{a_0} + 2R^2/a_0^2\right) \text{ where } \Psi(100) = (\pi a_0^3)^{-1/2} \exp(-r/a_0).$$

- (c) Write down the operators for L^2 and L_z in polar coordinates. Hence verify that $\Psi = A \sin \theta e^{i\phi}$, where A is a constant, is an eigenfunction of L^2 and L_z . Find the eigenvalues. 4+2+4

Section - II

(Atomic Physics)

Answer *any two* questions.

11. (a) In a Stern–Gerlach experiment, a beam of silver atoms moving with a velocity ‘ v ’ passes through an inhomogeneous magnetic field of gradient $\frac{\partial B}{\partial z}$ for a distance of ‘ l ’. After emerging from the magnetic field, they travel a distance ‘ b ’ before reaching the screen. What will be the magnitude of the splitting?
- (b) What is the g-factor for an atom with a single optical electron in $d_{3/2}$ level?
- (c) Consider the L-S coupling scheme for helium atom. Show that (i) $1s^1 2s^1$ configuration leads to the terms 1S_0 and 3S_1 while (ii) $1s^1 2p^1$ configuration leads to 1P_1 , 3P_0 , 3P_1 and 3P_2 . 4+2+(2+2)
12. (a) The spacing between the vibrational levels of CO molecule is 0.08 eV. Calculate the value of the force constant of the CO bond. Given that the masses of C and O atoms are 2.0×10^{-26} kg and 2.7×10^{-26} kg respectively. ($\hbar = 6.58 \times 10^{-16}$ eV sec)
- (b) Do hydrogen molecules give rise to pure vibration-rotation spectra? Justify your answer.
- (c) Pure rotational spectrum is almost always seen as absorption lines, and not as emission lines. Explain. 4+3+3
13. (a) Draw the energy level diagram for a four-level laser. Explain the requirement of each energy level. Why is a four-level laser preferred to a three-level laser?
- (b) In a He-Ne laser transition from $3S$ to $2P$ level gives a laser emission of wavelength 632.8 nm. If the $2P$ level has energy equal to $15.2 \times 10^{-19} J$, assuming no loss, calculate the pumping energy required.
- (c) Why do molecules show band spectra rather than line spectra? (2+3+1)+2+2